



## Epidemiological Surveillance of Respiratory Infections

### Weekly overview

**Week 28/2023 (10/7/2023 – 16/7/2023)**

#### Influenza-like Illness (ILI)

- ILI rate remained low

#### SARS-CoV2 virus - COVID-19 infection

- test positivity showed a small increase compared to the previous week
- the number of COVID-19 admissions (n=226) increased compared to the previous week and a 11% decrease compared to the average weekly number of new admissions during the previous 4 weeks
- the number of new intubations (n=2) decreased compared to the previous week and a 60% decrease compared to the average weekly number of new admissions during the previous 4 weeks
- the cumulative number of intubated patients with COVID-19 infection is 17
- 16 deaths were reported (median age: 84 years, range: 66-98)
- during the last weeks all sequenced samples were classified as Omicron sub-variants BA.2 and BA.5, with BA.2 being the dominant variant from week 9 onwards
- in week 24 the most frequent BA.2 sub-variants were XBB.1.5 (77%) followed by XBB.1.16 (16%)
- viral load surveillance in municipal wastewater showed an increase in SARS-CoV-2 virus circulation in 5 out of 10 areas participating in the network

#### Influenza virus

- all samples were tested negative for influenza virus
- no severe cases of laboratory-confirmed influenza admitted to ICU or deaths from laboratory-confirmed influenza were recorded in week 28/2023
- from week 40/2022 to week 28/2023, 68 cases were hospitalized in ICU and 26 deaths were reported
- during the same period, 374 samples positive for influenza viruses (sentinel samples and hospital samples) were detected in the two Influenza Reference Centers, of which 290 (77,5%) were type A and 84 (22,5%) were type B
- of the 288 type A viruses subtyped, 258 (90%) were classified as subtype A(H3N2) and 30 (10%) as subtype A(H1N1)pdm09

#### Respiratory syncytial virus – RSV

- all samples were tested negative for RSV