



## **Epidemiological surveillance in points of care for refugees/migrants Weekly Report**

**Week 27/2016 (04/07 to 10/07)**

### **SUMMARY**

In week 27/2016 (04/07 to 10/07) the system for epidemiological surveillance in points of care for refugees/migrants received data from 47 centres hosting refugees/migrants out of a total of 52 centres participating in the system (rate 90%).

During this week, the observed morbidity ranged within the expected limits, presenting only sporadic and short-term peaks of gastroenteritis in some centres. Two cases of the syndrome “jaundice of acute onset” were reported from one centre, which were further investigated and public health measures were taken. One case of the syndrome “meningitis and/or encephalitis” was reported, for which further investigation showed that it was not a meningitis case.

No case was recorded for the following syndromes/health conditions: [7] Malaria (with positive rapid test), [8] Suspected diphtheria (respiratory or cutaneous), [10] Paralytic manifestations of acute onset, [12] Haemorrhagic manifestations with fever, [13] Sepsis or shock (septic, of unknown aetiology), [14] Death of unknown aetiology (see table 1).

*This report is based on data recorded and sent by a large number of health professionals who provide health care, often under very difficult circumstances, in centres hosting refugees/migrants.*

*We thank all these professionals for their valuable contribution to monitoring morbidity among refugees/migrants, which is necessary for appropriate public health action.*

## A. Information from the system for epidemiological surveillance in points of care for refugees/migrants

During week 27/2016 (04/07 to 10/07), XX centres hosting refugees/migrants operated in Greece, from which 52 participate in the surveillance system and 47 sent data at least once during this week (rate 90%). The inclusion of centres in the surveillance system is ongoing.

Table 1 presents observed and expected morbidity data from centres hosting refugees/migrants.

Graphs 1 to 5 depict the time trend of the morbidity of the most frequent syndromes/health conditions monitored (for the rest of the syndromes, graphs are not shown, due to the small numbers of cases).

**Note.** Data presented here can be modified in the future, as delayed reports are included.

**Table 1:** Number of cases, proportional morbidity and statistical warning/alert signals by syndrome/health condition under surveillance, total of reporting centres hosting refugees/migrants, week 27/2016 (04/07 to 10/07).

| Syndrome                                           | No of cases | Obs. prop. morbidity | Exp. prop. morbidity | Z-score |
|----------------------------------------------------|-------------|----------------------|----------------------|---------|
| [1] Respiratory infection with fever               | 403         | 4,2                  | 4,2                  | -0,046  |
| [2] Gastroenteritis without blood in the stool     | 334         | 3,5                  | 3,3                  | 0,442   |
| [3] Bloody diarrhoea                               | 2           | 0,0                  | 0,0                  | -0,698  |
| [4] Rash with fever                                | 73          | 0,8                  | 0,6                  | 0,773   |
| [5] Suspected scabies                              | 44          | 0,5                  | 0,4                  | 0,310   |
| [6] Suspected pulmonary tuberculosis               | 2           | 0,0                  | 0,0                  | -0,359  |
| [7] Malaria with positive RDT                      | 0           | 0,0                  | 0,0                  | -0,207  |
| [8] Suspected diphtheria, respiratory or cutaneous | 0           | 0,0                  | 0,0                  | 0,000   |
| [9] Jaundice of acute onset                        | 3           | 0,0                  | 0,0                  | 0,987   |
| [10] Neurological manifestations of acute onset    | 0           | 0,0                  | 0,0                  | 0,000   |
| [11] Meningitis and/or encephalitis                | 1           | 0,0                  | 0,0                  | 0,723   |
| [12] Haemorrhagic manifestations with fever        | 0           | 0,0                  | 0,0                  | 0,000   |
| [13] Sepsis or shock (septic, of unknown etiology) | 0           | 0,0                  | 0,0                  | 0,000   |
| [14] Death of unknown etiology                     | 0           | 0,0                  | 0,0                  | 0,000   |


### Notes:

- Proportional morbidity: consultations for a given syndrome/health condition as a proportion of total consultations (for all causes).
- Observed proportional morbidity: refers to the week of the current report.
- Expected proportional morbidity: reflects the trend of the past 4 weeks.
- Z-score: difference between the observed and the expected proportional morbidity, expressed in number of standard deviations (Z-score > 2: observed proportional morbidity "statistically significantly" larger than the expected).

## Time trend of proportional morbidity of the monitored syndromes/health conditions with the highest frequency of occurrence in the total of participating centres hosting refugees/migrants


### 1. Respiratory infection with fever

Graph 1: Proportional morbidity of Respiratory infection with fever, based on reports from all camps in Greece



### 2. Gastroenteritis without blood in the stool

Graph 2: Proportional morbidity of Gastroenteritis without blood in the stool, based on reports from all camps in Greece




### 3. Bloody diarrhoea

Graph 3: Proportional morbidity of Bloody diarrhoea, based on reports from all camps in Greece

Point of care surveillance in refugee/migrant reception centres, Greece

Bloody diarrhoea




### 4. Rash with fever

Graph 4: Proportional morbidity of Rash with fever, based on reports from all camps in Greece


Point of care surveillance in refugee/migrant reception centres, Greece

Rash with fever



5. Suspected scabies

Graph 5: Proportional morbidity of Suspected scabies, based on reports from all camps in Greece



## **B. Additional information and public health measures taken**

During week 27 (04/07 – 10/07) three warning statistical signals of “gastroenteritis without blood in the stool” were investigated:

1. On 04/07 10 cases were reported from one center hosting refugees/migrants. All cases had mild symptoms, none was hospitalized and there was no further continuation of the outbreak.
2. On 07/07 13 cases were reported from one center hosting refugees/migrants. All cases had mild symptoms, none was hospitalized and there was no further continuation of the outbreak.
3. On 10/7 6 cases were reported from one center hosting refugees/migrants. All cases had mild symptoms, none was hospitalized and there was no further continuation of the outbreak.

In addition, a warning statistical signal for two reported cases of “jaundice with acute onset” was investigated in one center hosting refugees/migrants. After laboratory investigation the cases were confirmed as hepatitis A. Vaccination of close contacts was scheduled in collaboration with the Hellenic Red Cross.

During this week, 38 cases of chickenpox were reported from the total of the refugees/migrants centers in the country, the majority of whom were children (35 cases aged up to 14 years old). Chickenpox is usually a mild disease in childhood; populations in which childhood vaccination has not been introduced (i.e. populations from the countries of origin of refugees/migrants hosted in the centers) experience chickenpox as a “childhood disease”, with periodic outbreaks. The HCDCP has issued instructions for the management of chickenpox cases in centers hosting refugees/migrants ([www.keelpno.gr/](http://www.keelpno.gr/)).

## C. Methods

Public health surveillance in points of care for refugees/migrants operates in its present form from 16 May 2016 (and as a pilot from April 2016) with daily collection of epidemiological data for selected syndromes/health conditions that are important from a public health point of view. The 14 syndrome or health conditions under surveillance are shown in Table 1.

Data recorded refer to consultations for each syndrome/condition under surveillance in primary health care facilities in refugee/migrant reception centres (RMRC). For syndromes #1 to 5, which have the highest incidence, cumulative data are collected (i.e. number of consultations without any additional information), while for syndromes #6 to 14 some important individual-level information is also collected for patients. In addition, individual-level information is collected for cases with clinical suspicion of measles, rubella, mumps and varicella.

Data are sent daily to the Department of Surveillance and Intervention of KEELPNO by doctors, nurses and other health professionals from services and NGOs staffing primary care facilities in RMRCs.

Data for a given 24-hour period are analysed on the next day and proportional morbidity indices are calculated (consultations for each syndrome/condition under surveillance as a percentage of the total number of consultations, i.e. the number of consultations for all causes). This analysis is carried out for all RMRCs in the country participating in the system as a whole, and for each RMRC separately. Moreover, weekly data are also analysed (see Table 1).

The index of proportional morbidity (observed proportional morbidity) is compared with the expected proportional morbidity, which is calculated using a statistical model\*. The expected proportional morbidity reflects the trend of the preceding 4 weeks; determining the warning threshold takes into account the dispersion of the daily values of proportional morbidity during the whole period since 16/05/2016. Observed proportional morbidity higher than the expected by more than 2 standard deviations ( $Z\text{-score} > 2$ ) is equivalent to a "warning signal". A "warning signal" that appears for at least two consecutive days is equivalent to an "alert signal". The signals are evaluated in terms of importance for public health and, if necessary, they are further investigated, and –if considered appropriate– public health action is organized.

---

\* The expected proportional morbidity is calculated using a quasi-Poisson regression model (Farrington et al, 1996). Long-term trends are incorporated in the model using natural cubic splines (with knots every 4 weeks), rejecting outlier values ( $Z\text{-score} > 3$ ) and –if considered necessary– values corresponding to a confirmed outbreak.