

## Weekly Epidemiological Report for West Nile Virus disease, Greece, 2017 - 26 July 2017 -

This weekly epidemiological report aims to present an overview of the reported cases and public health response to West Nile Virus (WNV) in Greece for transmission period 2017.

Data presented in this report are derived from the notifications of laboratory diagnosed human cases of WNV infection sent to the Hellenic Center for Disease Control and Prevention (HCDCP-KEELPNO) by the treating physicians and from the daily communication with diagnostic laboratories: i) the Reference Laboratory for Arboviruses, Aristotelian University of Thessaloniki, ii) the Department of Microbiology, School of Medicine, University of Athens, iii) the Hellenic Pasteur Institute. The Department of Epidemiological Surveillance and Intervention of the HCDCP undertakes a verification procedure through communication with the treating physicians and the patients, as necessary.

From the beginning of 2017 until 25/07/2017 (16:00), five (5) laboratory diagnosed cases of WNV infection have been reported to KEELPNO, all of which presented with neuro-invasive disease (WNND, encephalitis and/or meningitis and/or acute flaccid paralysis) ([Table 1](#)).


**Table 1. Number of reported cases of WNV disease, Greece, period 2017, until 25/07/2017 (16:00)**

	Number of cases <b>with</b> central nervous system (CNS) manifestations <sup>[1]</sup>	Number of cases <b>without</b> CNS manifestations	<b>Total number of cases</b>	<b>Number of deaths</b> <sup>[2]</sup>
Number of WNV cases and deaths	5	0	5	0

1. Refers mainly to encephalitis, aseptic meningitis and meningoencephalitis cases
2. The number of deaths is included in the total number of cases

[Figure 1](#) shows the reported WNND cases by week of symptom onset. The first diagnosed case of WNV infection for transmission period 2017 reported onset of symptoms on 20<sup>th</sup> June 2017 (wk 25/2017).

**Figure 1. Number of laboratory diagnosed WNND cases by week of symptom onset, Greece, 2017, until 25/07/2017, 16.00 (n=5)\***



\* Each orange box represents one laboratory diagnosed case of WNND reported to KEELPNO in transmission period 2017.

The median age of WNND cases is 63 years (27-73 years).

[Table 3](#) show the geographic distribution of the notified cases with laboratory diagnosed WNV disease at the level of suspected Municipalities of exposure. The patient’s suspected place of exposure is a rough indicator of the area of WNV circulation.

According to a serosurvey conducted in 2010 by the HCDCP and the National School of Public Health, at the epicentre of the 2010 WNV outbreak in Central Macedonia, WNND disease develops in 1:140 infected persons.

**Table 3. Reported cases with laboratory diagnosed WNV disease (with and without WNND) by suspected Municipality of exposure, Greece, transmission period 2017, until 25/07/2017, (16:00)**

Regional Unit	Suspected Municipality of exposure	Number of cases with WNND	Incidence of WNND per 100,000 population *	Number of West Nile Fever cases
Argolis	Argos-Mykines	2	4.76	0
	Nafplio	3	8.99	0
<b>Total, Greece</b>		<b>5</b>	<b>0.05</b>	<b>0</b>

\* Calculations based on 2011 census data (Hellenic Statistical Authority).

### **PUBLIC HEALTH MEASURES SUPPORTED BY THE HCDCP - 2017**

The following public health measures have been implemented by the HCDCP and other involved stakeholders:

#### **I. Enhanced surveillance for encephalitis and WNV disease in humans:**

- **Awareness raising of physicians** about the WNV infection: Testing for West Nile virus infection in suspected cases (such as cases with encephalitis, aseptic meningitis, acute flaccid paralysis, fever of undetermined etiology) is recommended. The HCDCP provided guidelines for the recognition and diagnosis of WNV disease and the recommended laboratory investigation (mailings and website [www.keelpno.gr](http://www.keelpno.gr)).
- **Daily communication and information exchange with laboratories** conducting diagnostic testing for WNV.
- **Enhancing laboratory diagnosis** of suspected cases, by supporting specialised diagnostic laboratories.
- **Case investigation:** The Department of Epidemiological Surveillance and Intervention of HCDCP undertakes the investigation of every reported WNV case within 24 hours after diagnosis, in order to determine the likely place of exposure, the risk factors and the severity of the disease.
- **Daily update of stakeholders** on the diagnosed cases (Ministry of Health, Ministry of Rural Development and Food, Regions, Directorate of Public Health and Social Welfare, National Centre for Blood Donations).

II. **Communication and health promotion activities for the public:** Distribution of leaflets with educational material for the public regarding the recommended protective measures against mosquito bites in collaboration with local authorities; educational material in the HCDCP website. In 2017, information material (leaflets, posters) has been created and distributed to the Regional authorities, according to their needs (upon request). In addition, HCDCP sends leaflets and posters immediately in areas where cases are recorded.

III. **Coordination of an intersectional Working Group (WG) on the definition of affected areas by vector borne diseases.** This WG, under the MoH Committee for the Prevention and Management of Tropical Diseases, considers all available entomological and epidemiological data and decides on the characterization of affected areas assisting the implementation of blood safety measures. The list of affected areas is published on our website ([www.keelpno.gr](http://www.keelpno.gr)) and updated regularly. These are used by

**HELLENIC CENTER FOR DISEASE CONTROL AND PREVENTION (HCDCP)**

Office for Vector Borne Diseases

Department for Epidemiological Surveillance and Intervention

210 8899 052, 210 8899 071, 210 8899 072

the National Centre for Blood Donation to issue guidance on blood safety. In addition, the Hellenic Centre for Coordinating Haemovigilance has issued guidance for the haemovigilance competent authorities.

**IV.** Collaboration and exchange of information with the **Ministry of Rural Development and Food** regarding the West Nile virus infection in equids.

**V. Vector surveillance and control activities:**

- **Raising awareness and guidance to Regional Authorities:** HCDCP communicates regularly (workshops, meetings, letters) with all Regional Authorities in Greece recommending the timely planning, organization and implementation of integrated vector control programmes. In 2017, HCDCP send relevant awareness letters in January 2017 (with a brief guide to the key steps to achieve timely implementation of the vector control program). In addition, at the initiative of the Secretary General of Public Health of the Ministry of Health and the Association of the Regions of Greece, national and regional public health services have held working group meetings regarding the preventive actions for the mosquito-borne diseases.

Following the first case of the WNV infection, a working group meeting was held among the regional and local authorities and the HCDCP, regarding the preventive and response activities (intensifying the mosquito control programs and raising awareness of the local population).

- **Monitoring of the vector control programmes' stage** at each Region/ Regional Unit.
- **Entomological surveillance:** The HCDCP, in collaboration with the Department of Parasitology, Entomology and Tropical Diseases of the National School of Public Health (NSPH), the Benaki Phytopathological Institute, the MALWEST project (2012-2014), Universities, Regions, local authorities and subcontractors of the local mosquito control programmes has implemented, participated or coordinated -from 2010 to 2015- active vector surveillance programme. For the 2017 transmission period, vector surveillance is organized to be performed in various areas of the country. HCDCP recommends to local authorities to perform vector surveillance, especially in the affected areas.
- **Communication with international public health stakeholders:** Frequent communication and weekly information exchange with ECDC (real-time reporting of the diagnosed cases in TESSy).

**CONCLUSIONS**

During the period 2010-2014 cases of West Nile virus infection were recorded in humans in various regions of the country, while virus circulation was recorded in almost all regions. Although years 2015-2016 no cases of the infection was recorded in humans in Greece, given the complex epidemiology and the unpredictable virus circulation, the recurrence of WNV infection cases in the country was considered likely and expected.

Until 25/07/2017, human WNV cases were recorded from the Regional Unit of Argolis, new area of virus circulation.

Epidemiological surveillance of disease, systematic and early implementation of mosquito control programs and personal protective measures against mosquito bites are the most appropriate measures for the control of the disease.

Since the areas of virus circulation during the current period cannot be predicted, personal protective measures against mosquitoes are strongly encouraged all over the country.